

DAFTAR PUSTAKA

1. Sugiharto A. Faktor-faktor risiko hipertensi Grade II pada masyarakat (Studi Kasus di Kabupaten Karanganyar). Diponegoro University Institutional Repository. 2007;.
2. World Health Organization. Take Control High Blood Pressure Global and Regional Overview [Internet]. Geneva: WHO; 2013 [cited 8 August 2018]. Available from: http://www.searo.who.int/entity/world_health_day/leaflet_burden_hbp_whd2013.pdf?ua=1
3. Kementerian Kesehatan Republik Indonesia. Hasil Utama Riskesdas 2018 [Internet]. Jakarta: Kementerian Kesehatan Republik Indonesia; 2018 [cited 2018 November 9]. Available from: <http://depkes.go.id>
4. Artiyaningrum B. Faktor-faktor yang berhubungan dengan kejadian hipertensi tidak terkontrol pada penderita yang melakukan pemeriksaan rutin di puskesmas kedungmundu kota Semarang tahun 2014. Jurnal Ilmu Kesehatan Masyarakat Fakultas Ilmu Keolahragaan Universitas Negeri Semarang; 2015.
5. Nuraini B. Risk Factors of Hypertension. J Majority. 2015;4(5).
6. Andrea G. Korelasi Derajat Hipertensi dengan Stadium Penyakit Ginjal Kronik di RSUP Dr. Kariadi Semarang periode 2008-2012. Diponegoro University Institutional Repository. 2013;.
7. Hasanah M, Widodo D, Widiani E. Hubungan Obesitas dengan Hipertensi pada Masyarakat di Wilayah RW 13 Dusun Mojosari Desa Ngenep Kecamatan Karangploso. Nursing News. 2016;1(2).
8. Nuertey B, Alhassan A, Nuertey A, Mensah I, Adongo V, Kabutey C et al. Prevalence of obesity and overweight and its associated factors among registered pensioners in Ghana; a cross sectional studies. BMC Obesity. 2017;4(1).
9. Ramadhani R. Hubungan Obesitas dengan Kejadian Hipertensi pada Masyarakat Etnik Minangkabau di Kota Padang. Bagian Gizi Fakultas Kedokteran Universitas Andalas. 2018;.

10. National Health and Nutrition Examination Survey, Centers for Disease Control and Prevention (CDC). Anthropometry Procedures Manual. United States: Centers for Disease Control and Prevention; 2007 [cited 2018 November 19]. Available from: https://www.cdc.gov/nchs/data/nhanes/nhanes_07_08/manual_an.pdf
11. Li W-C, Chen I-C, Chang Y-C, Loke S-S, Wang S-H, Hsiao K-Y. Waist-to-height ratio, waist circumference, and body mass index as indices of cardiometabolic risk among 36,642 Taiwanese adults. *European Journal of Nutrition* [Internet]. 2011 Dec 11;52(1):57–65. Available from: <http://dx.doi.org/10.1007/s00394-011-0286-0>
12. Ashwell M, Gibson S. Waist-to-height ratio as an indicator of ‘early health risk’: simpler and more predictive than using a ‘matrix’ based on BMI and waist circumference. *BMJ Open* 2016;6:e010159. doi:10.1136/bmjopen-2015-010159
13. Kartika TA, Utari DM. Faktor Dominan Kejadian Pre Hipertensi dan Hipertensi pada Petugas Satpam Laki-Laki Universitas Indonesia Tahun 2014. Departemen Ilmu Gizi, Fakultas Kesehatan Masyarakat, Universitas Indonesia. 2014.
14. Soenarta A, Mumpuni A. PEDOMAN TATALAKSANA HIPERTENSI PADA PENYAKIT KARDIOVASKULAR. 1st ed. Perhimpunan Dokter Spesialis Kardiovaskular Indonesia; 2019.
15. Kurniasih F, Meila O, Tasriyanti F. Pemantauan Terapi Obat pada Pasien Hipertensi di Bangsal Flamboyan Rumah Sakit Tk.II dr. AK Gani Palembang Periode 01 Maret – 26 April 2017. *Social Clinical Pharmacy Indonesia Journal Universitas 17 Agustus 1945 Jakarta*. 2017;2(1).
16. Qureshi AI. Acute Hypertensive Response in Patients With Stroke. *Circulation*. 2008Aug;118(2):176–87.
17. Infodatin, depkes. Hipertensi [Internet]. Indonesia: Departemen Kesehatan; 2014 [cited 11 November 2018]: Available from: <http://www.depkes.go.id/download.php?file=download/pusdatin/infodatin/infodatin-hipertensi.pdf>
18. Chobanian AV. The Seventh Report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood

- Pressure<SUBTITLE>The JNC 7 Report</SUBTITLE>. JAMA [Internet]. 2003 May 21;289(19):2560. Available from: <http://dx.doi.org/10.1001/jama.289.19.2560>
19. World Health Organization. Raised blood pressure [Internet]. Geneva: World Health Organization; 2018 [cited 18 September 2018]. Available from: https://www.who.int/gho/ncd/risk_factors/blood_pressure_prevalence_text/en/
 20. Ashwell M, Lejeune S. Ratio of waist circumference to height may be better indicator of need for weight management. BMJ [Internet]. 1996 Feb 10;312(7027):377–377. Available from: <http://dx.doi.org/10.1136/bmj.312.7027.377>
 21. Gibson, R. Principles of nutritional assessment. New York: Oxford University Press. 2005.
 22. Schneider, H. J., Klotsche, J., Silber, S., Stalla, G. K., & Wittchen, H.-U. Measuring Abdominal Obesity: Effects of Height on Distribution of Cardiometabolic Risk Factors Risk Using Waist Circumference and Waist-to-Height Ratio. 2010. Diabetes Care, 34(1), e7–e7. <https://doi.org/10.2337/dc10-1794>
 23. Tawfik, H. Waist height ratio and waist circumference in relation to hypertension, Framingham risk score in hospitalized elderly Egyptians. The Egyptian Heart Journal, 2018. 70(3), pp.213-216.
 24. Indonesia Fitness Trainer Association. Cara Mengukur Tinggi dan Berat Badan - Indonesia Fitness Trainer Association [Internet]. Indonesia: Indonesia Fitness trainer Association; 2018 [cited 27 November 2018]. Available from: <https://www.apki.or.id/cara-mengukur-tinggi-dan-berat-badan/>
 25. World Health Organization. Waist Circumference and Waist-Hip Ratio: report of a WHO Expert Consultation [Internet]. Geneva: WHO Document Production of Services; 2008 [cited 2 December 2018]. Available from: http://whqlibdoc.who.int/publications/2011/9789241501491_eng.pdf

26. Sherwood L. Introduction to human physiology. 8th ed. Pacific Grove, Calif.: Brooks/Cole; 2013. Chapter 10, Blood Vessels and Blood Pressure; p.368-369.
27. Grasgruber P, Sebera M, Hrazdíra E, Cacek J, Kalina T. Major correlates of male height: A study of 105 countries. *Economics & Human Biology* [Internet]. 2016 May;21:172–95.
28. Bourgeois B, Watts K, Thomas DM, Carmichael O, Hu FB, Heo M, et al. Associations between height and blood pressure in the United States population. *Medicine* [Internet]. 2017 Dec;96(50):e9233.
29. Browning LM, Hsieh SD, Ashwell M. A systematic review of waist-to-height ratio as a screening tool for the prediction of cardiovascular disease and diabetes: 0.5 could be a suitable global boundary value. *Nutr Res Rev*. 2010;23:247–269
30. Sari M, Lipoeto N, Herman R. Hubungan Lingkar Abdomen (Lingkar Perut) dengan Tekanan Darah. *Jurnal Kesehatan Andalas*. 2016;5(2):456-461
31. V Roth J. Taller people should have Higher BMI's and Blood Pressure Measurements as their Normal. *Biomedical Journal of Scientific & Technical Research* [Internet]. 2018 Jul 10;6(4).
32. Ashwell M, Hsieh SD 2005 Six reasons why the waist-to-height ratio is a rapid and effective global indicator for health risks of obesity and how its use could simplify the international public health message on obesity. *Int J Food Sci Nutr* 56:303–307