

ISSN-0976-0245 (Print) • ISSN-0976-5506 (Electronic)

Volume 9

Number 12

December 2018



Indian Journal of Public Health Research & Development

An International Journal

SCOPUS IJPHRD CITATION SCORE

Indian Journal of Public Health Research and Development

Scopus coverage years: from 2010 to 2017 Publisher:

R.K. Sharma, Institute of Medico-Legal Publications


ISSN:0976-0245E-ISSN: 0976-5506 Subject area: Medicine:

Public Health, Environmental and Occupational Health

CiteScore 2015- 0.02

SJR 2015-0.105

SNIP 2015- 0.034



Website:

www.ijphrd.com

Indian Journal of Public Health Research & Development

EXECUTIVE EDITOR

Vidya Surwade

Associate Professor, Dr Baba Saheb Ambedkar, Medical College & Hospital, Rohinee, Delhi

INTERNATIONAL EDITORIAL ADVISORY BOARD

1. **Dr. Abdul Rashid Khan B. Md. Jagar Din**, (Associate Professor)
Department of Public Health Medicine, Penang Medical College, Penang, Malaysia
2. **Dr. V Kumar** (Consulting Physician)
Mount View Hospital, Las Vegas, USA
3. **Basheer A. Al-Sum**,
Botany and Microbiology Deptt, College of Science, King Saud University,
Riyadh, Saudi Arabia
4. **Dr. Ch Vijay Kumar** (Associate Professor)
Public Health and Community Medicine, University of Buraimi, Oman
5. **Dr. VMC Ramaswamy** (Senior Lecturer)
Department of Pathology, International Medical University, Bukit Jalil, Kuala Lumpur
6. **Kartavya J. Vyas** (Clinical Researcher)
Department of Deployment Health Research,
Naval Health Research Center, San Diego, CA (USA)
7. **Prof. PK Pokharel** (Community Medicine)
BP Koirala Institute of Health Sciences, Nepal
8. **Sajjad Salim Issa AL-Musawi**, Supervisor for Student of the Arab
Board in Family Medicine
9. **Prof. Dr. Ayad F. Alkaim**, Professor of Surface Chemistry and
Applications of Nano-materials, Babylon University, Iraq

NATIONAL SCIENTIFIC COMMITTEE

1. **Dr. Anju Ade** (Associate Professor)
Navodaya Medical College, Raichur, Karnataka
2. **Dr. E. Venkata Rao** (Associate Professor) Community Medicine,
Institute of Medical Sciences & SUM Hospital, Bhubaneswar, Orissa.
3. **Dr. Amit K. Singh** (Associate Professor) Community Medicine,
VCSG Govt. Medical College, Srinagar – Garhwal, Uttarakhand
4. **Dr. R G Viveki** (Professor & Head) Community Medicine,
Belgaum Institute of Medical Sciences, Belgaum, Karnataka
5. **Dr. Santosh Kumar Mulage** (Assistant Professor)
Anatomy, Raichur Institute of Medical Sciences Raichur(RIMS), Karnataka
6. **Dr. Gouri Ku. Padhy** (Associate Professor) Community and Family
Medicine, All India Institute of Medical Sciences, Raipur
7. **Dr. Ritu Goyal** (Associate Professor)
Anaesthesia, Sarswathi Institute of Medical Sciences, Panchsheel Nagar
8. **Dr. Anand Kalaskar** (Associate Professor)
Microbiology, Prathima Institute of Medical Sciences, AP
9. **Dr. Md. Amirul Hassan** (Associate Professor)
Community Medicine, Government Medical College, Ambedkar Nagar, UP
10. **Dr. N. Girish** (Associate Professor) Microbiology, VIMS&RC, Bangalore
11. **Dr. BR Hungund** (Associate Professor) Pathology, JNMC, Belgaum.
12. **Dr. Sartaj Ahmad** (Assistant Professor),
Medical Sociology, Department of Community Medicine, Swami Vivekananda Subharti
University, Meerut, Uttar Pradesh, India
13. **Dr Sumeeta Soni** (Associate Professor)
Microbiology Department, B.J. Medical College, Ahmedabad, Gujarat, India

NATIONAL EDITORIAL ADVISORY BOARD

1. **Prof. Sushanta Kumar Mishra** (Community Medicine)
GSL Medical College – Rajahmundry, Karnataka
2. **Prof. D.K. Srivastava** (Medical Biochemistry)
Jamia Hamdard Medical College, New Delhi
3. **Prof. M Sriharibabu** (General Medicine) GSL Medical College, Rajahmundry,
Andhra Pradesh
4. **Prof. Pankaj Datta** (Principal & Prosthodontist)
Indraprastha Dental College, Ghaziabad

NATIONAL EDITORIAL ADVISORY BOARD

5. **Prof. Samarendra Mahapatro** (Pediatrician)
Hi-Tech Medical College, Bhubaneswar, Orissa
6. **Dr. Abhiruchi Galhotra** (Additional Professor) Community and Family
Medicine, All India Institute of Medical Sciences, Raipur
7. **Prof. Deepti Pruthvi** (Pathologist) SS Institute of Medical Sciences &
Research Center, Davangere, Karnataka
8. **Prof. G S Meena** (Director Professor)
Maulana Azad Medical College, New Delhi
9. **Prof. Pradeep Khanna** (Community Medicine)
Post Graduate Institute of Medical Sciences, Rohtak, Haryana
10. **Dr. Sunil Mehra** (Paediatrician & Executive Director)
MAMTA Health Institute of Mother & Child, New Delhi
11. **Dr. Shailendra Handu**, Associate Professor, Pharma, DM (Pharma, PGI
Chandigarh)
12. **Dr. A.C. Dhariwal**: Directorate of National Vector Borne Disease
Control Programme, Dte. DGHS, Ministry of Health Services, Govt. of
India, Delhi

Print-ISSN: 0976-0245-Electronic-ISSN: 0976-5506, Frequency: Monthly

Indian Journal of Public Health Research & Development is a double blind peer reviewed international journal. It deals with all aspects of Public Health including Community Medicine, Public Health, Epidemiology, Occupational Health, Environmental Hazards, Clinical Research, and Public Health Laws and covers all medical specialties concerned with research and development for the masses. The journal strongly encourages reports of research carried out within Indian continent and South East Asia.

The journal has been assigned International Standards Serial Number (ISSN) and is indexed with Index Copernicus (Poland). It is also brought to notice that the journal is being covered by many international databases. The journal is covered by EBSCO (USA), Embase, EMCare & Scopus database. The journal is now part of DST, CSIR, and UGC consortia.

Website : www.ijphrd.com

©All right reserved. The views and opinions expressed are of the authors and not of the Indian Journal of Public Health Research & Development. The journal does not guarantee directly or indirectly the quality or efficacy of any product or service featured in the advertisement in the journal, which are purely commercial.

Editor

Dr. R.K. Sharma
Institute of Medico-legal Publications
Logix Office Tower, Unit No. 1704, Logix City Centre Mall,
Sector- 32, Noida - 201 301 (Uttar Pradesh)

Printed, published and owned by

Dr. R.K. Sharma
Institute of Medico-legal Publications
Logix Office Tower, Unit No. 1704, Logix City Centre Mall,
Sector- 32, Noida - 201 301 (Uttar Pradesh)

Published at

Institute of Medico-legal Publications
Logix Office Tower, Unit No. 1704, Logix City Centre Mall,
Sector- 32, Noida - 201 301 (Uttar Pradesh)



Indian Journal of Public Health Research & Development

www.ijphrd.com

CONTENTS

Volume 9, Number 12

December 2018

1. The Etiological Profile of Seizures in Children in a Tertiary Care Hospital, Hapur, Uttar Pradesh 01
Shweta Singh, M Agrawal, Yogesh Kumar Goel, Dayachand Verma
2. Does Pregnant Mother are aware about PMTCT of HIV? What is their Attitude and Do they Practice the Measures on PMTCT of HIV?: A Narrative Review 07
Neethu Sabu, N V Muninarayanappa, Kavitha Mole P J
3. Recent Resting Habit of Adult *Phlebotomus argentipes* the Vector of *Visceral leishmaniasis* in a Kala Azar Endemic Foci of Bihar India 12
Chandrima Das, Shilpa Raj, A K Mukhopadhyay
4. Substance Use Disorder - Vital Hurdle in Sustainable Development of Nigeria 18
Jummai Fatima Muhammad, Malavika Bhattacharya
5. A Study of Urinary Uric Acid/Creatinine Ratio as an Additional Marker of Birth Asphyxia 23
Renu Yadav, Sangeeta Singhal, Gagan Agarwal
6. Incidence, Prevalence and Mortality of HIV/AIDS across Different Levels of Human Development Index: A Global Perspective 28
Ravi Prakash Jha, Kritika Bhattacharyya, Rabindra Nath Mishra, Akash Mishra
7. Study to Assess the Social and Economic Impact of Alcohol in a Peri-Urban Area of Tamil Nadu 34
Shankar S, Raghuram V, Elango S
8. Economics of Sustainability – A Theoretical Perspective 40
S N Sugumar, S Balasekaran, S Chandrachud
9. Factors Affecting Neonatal Mortality and Morbidity - An Epidemiological Study 45
Tapan Pattanaik, Ratan Kumar Das, K Trimal Subudhi, Mahesh Chandra Sahu
10. Students' Perception and Attitude on Education Curriculum and System in an Indian Dental School 50
Madhura Sen, Kundabala M
11. Perception of Anganwadi Services in Urban ICDS Blocks in Kozhikode Corporation –A Cross -Sectional Study 55
Sivakumar D, Asma Rahim, Deepika

121. Joint Effect of Obesity and Cigarette Smoking Against Hypertension Stage 1 among Men Adults: Finding from the Indonesian Family Life Survey-5 673
Aprizal Satria Hanafi, Nurhayati A Prihartono
122. Comparison of Various Obesity Indices against the Occurrence of Stage 1 Hypertension in Indonesia 679
Anggun Pratiwi, Nurhayati A Prihartono
123. The Status of Lipid Peroxidation System in Erysipelas Patients on the Background of Diabetes Mellitus 686
Madina Marjokhova, Marina Afashagova, Asiat Marjokhova, Maryana Nagoeva, Marina Ivanova, Marina Nalchikova, Zarema Tagirova
124. Environmental Health Risk due to Exposure to Lead in Batteries Smelter Industry- Cinangka Village, Bogor, Indonesia 691
Ladyka Viola, Haryoto Kusnoputranto, Bambang Wispriyono
125. Evaluation of Bone Mineral Density in Adult Epileptic Patients Treated with Valproate 698
Mohammad A.S. Kamil, Aqeel K. Hatem, Mustafa Easa
126. Policy and Role Analysis Integrated Health Education Centers for Non-Communicable Diseases Toward the Prevention and Controlling of Hypertension 705
Yandrizal, Rizanda Machmud, Melinda Noer, Hardisman, Ekowati Rahajeng, Desri Suryani, P.A Kodrat Pramudho
127. Association of Apolipoprotein E Polymorphism with Cognitive Functions in Elderly 711
N. Saelan Tadjudin, Andi Jayalangkara Tanra, Andi Asadul Islam, Suryani As'ad, Muh. Nasrum Massi, Ilhamjaya Patellongi, Prihantono Prihantono
128. Designing Instrument for Early Stimulation, Detection, and Intervention for Growth and Development of Children based on Android System 717
Heru Santoso Wahito Nugroho, Sunarto, Budi Joko Santosa
129. Effectiveness of the Training Model to Increase the Knowledge, Attitude, and Practice of Traditional Birth Attendants about Early Initiation of Breastfeeding in Bogor District, Indonesia 723
Evi Martha, Hadi Pratomo, Siti N. Qomariyah

Association of Apolipoprotein E Polymorphism with Cognitive Functions in Elderly

N. Saelan Tadjudin¹, Andi Jayalangkara Tanra², Andi Asadul Islam³, Suryani As'ad⁴, Muh. Nasrum Massi⁵, Ilhamjaya Patellongi⁶, Prihantono Prihantono³

¹Department of Psychiatry, Faculty of Medicine, Tarumanagara University, Jakarta, Indonesia, ²Department of Psychiatry, ³Department of Surgery, ⁴Department of Nutrition, ⁵Department of Microbiology, ⁶Department of Physiology, Faculty of Medicine, Hasanuddin University, Makassar, Indonesia

ABSTRACT

Objective: This study aims to examine the association between apolipoprotein E polymorphism (APOE) and cognitive functions test using Montreal Cognitive Assessment- Indonesian version 2012 (MoCA-Ina) in elderly.

Method: A cross-sectional study of 112 elderly participants from two nursing homes in Jakarta. APOE gene polymorphism was analyzed using polymerase chain reaction-restriction fragment length polymorphism (PCR-RFLP).

Results: There are four genotypes: E2/E2 10(8.9%), E3/E3 38(33.9%), E3/E4 46(41.1%) and E4/E4 18(16.1%). The percentage of E2 allele 20(8.9%), E3 122(54.5%), E4 82(36.6%). The cognitive function test, MoCA-Ina ≥ 26 : 34(40%) participants with genotype E2/E2 3(8.8%), E3/E3 7(20.6%), E3/E4 16(47.1%) and E4/E4 8(23.5%). MoCA-Ina < 26 : 51(60%) 51 participants with genotype E2/E2 7(13.7%), E3/E3 14(27.5%), E3/E4 22(43.1%) and E4/E4 8(15.7%). A significant relationship between APOE gene polymorphism with MoCA-Ina-abstract: E3/E4 vs. E3/E3 $p = 0.019$, E4/E4 vs. E3/E3 $p = 0.028$, a significant relationship between APOE gene polymorphism with MoCA-Ina-language: E3/E4 vs. E2/E2 $p = 0.03$, E4/E4 vs. E2/E2 $p = 0.029$, and a significant relationship between APOE gene polymorphism with naming: E3/E4 vs. E2/E2 $p = 0.048$.

Conclusion: High percentage of APOE allele E4 may be one of the causes of high cognitive function decline. APOE gene polymorphisms associated with the decreased cognitive function; abstraction, language, and naming. Allele E4 may be a potential factor for cognitive function decline.

Keywords: MoCA-Ina, Elderly, polymorphism, APOE, gene.

INTRODUCTION

Dementia is a complex neurodegenerative disease in the elderly with multifactorial causes and characterized by various cognitive abnormalities, which could interfere with daily activities. ^(1,2) Alzheimer's Disease (AD) is one of the most common types of dementia and associated with the aging process. AD results in a gradual decline in cognitive functions and daily activity due to degeneration processes that damage the brain structures and neurotransmitters. ^(1,3) In the AD, the patient's ability to remember, understand, communicate and comprehend causality decreases gradually, which ultimately lead

to death. ^(1,3-5) Many factors can clinically affect the progress of the damage, including brain atrophy^(1,6-8), vascular factor^(1,9) and immunological factor.^(1,10) Currently 35 million people have dementia worldwide, expected to reach 65 million by 2030 and 113 million by 2050. ⁽⁵⁾ The prevalence of dementia in Indonesian older adults aged 65 years is 5% and increases to 20% at aged above 85. ⁽¹¹⁾

A genetic factor that is known to play a role in the development of AD is the APOE gene polymorphism, lies on chromosome 19 and has three alleles: E2, E3, E4 and six genotype variations: E2/E2, E2/E3, E2/E4, E3/

E3, E3/E4 and E4/E4.^(12, 13) The allele E4 is associated with an increased risk of the AD (14-16), whereas the allele E2 appears to be a protective factor.^(5, 17) Having APOE genotype E4/E4 not only increases the risk of AD but also affects the cognitive symptoms and neuroanatomical forms of the disease. E4 carriers tend to have worse memory decline compared to non-carriers, while the APOE E4 non-carrier tend to have more difficulty maintaining attention, working memory, executive functions, and lexical access.⁽¹⁴⁾

This genetic factor is stronger in the European population⁽¹⁸⁾ and weaker in the African, American and Hispanic community.^(5, 19) Recent studies show that the effects of APOE E4 allele on cognitive functions depend on the age and level of education of the individuals.⁽²⁰⁾ It is still debatable whether APOE E4 is related to cognitive functions or cognitive impairment.⁽²⁰⁾ With polymerase chain reaction-restriction fragment length polymorphism, the association between APOE E4 polymorphism and cognitive function was analyzed.

MATERIALS AND METHOD

Collection of Samples

Participants of this a cross-sectional was 112 elderly aged between 60 - 90 years, consisting of 99 women and 13 men from two nursing homes in Jakarta. The DNA was analyzed to find out APOE genotypes. Cognitive functions examination was carried out on 85 female participants using MoCA-Ina questionnaire and its items (visuospatial, naming, attention, language, abstraction, memory delay, orientation) from the neurology department, University of Indonesia. MoCA-Ina scores divided into normal cognitive function ≥ 26 and cognitive impairment < 26 .

PCR-RFLP Analysis

Blood samples obtained from a peripheral vein using 3 ml vacutainer tubes containing EDTA. The blood samples were centrifuged for 30 minutes at 3000 rpm and separated into serum from sediment. The sediment Examined for APOE genes. DNA is extracted from blood samples using a DNA Extraction Kit (Bioline) according to the manufacturer's instructions.

DNA extraction

200 μ l of the blood sample inserted into a 1.5 ml sterile microcentrifuge tube. 20 μ l Proteinase K then

added into the tube. The solution is homogenized by pipetting and then incubated at 60 ° C for 5 minutes. Add 200 μ l Buffer GSB (Geneid) and vortex the tube. Incubate the tube again at the same temperature for 2 minutes. Add ethanol absolute (96%) and vortex for 10 seconds. Pour the mixture into the spin column and centrifuge at 14,000 xg for 1 minute. Remove the collection tube under the spin column and replace it with a new one. Add 400 μ l of buffer W1 and centrifuge for 30 seconds with the same speed. Discard the liquid that is on the collection tube. Add 600 μ l of wash buffer (Geneid), centrifuge the mixture for 30 seconds, and dispose of liquid in the collection tube. Centrifuge again for 3 minutes. Remove the collection tube and place a sterile microcentrifuge below the spin column. Next, add 100 μ l of Elution buffer. Let the mixture stand for 3 minutes and then centrifuge with the same speed for 30 seconds. The fluid containing the DNA stored in a microcentrifuge tube at -20 ° C and will be used as template DNA for PCR.

The genotype of the APOE gene

The extracted DNA added into PCR Mix, which consists of 5 μ L 5X buffer, 2 μ L MgCl₂, 1 μ L dNTP mix, 1 μ L APOE-F Primer (5'-ACA GAA TTC GCC CCG GCC TGG TAC AC-3'), 1 μ L Primary APOE-R (ApoE-R: 5'-TAA GCT TGG CAC GGC TGT CCA AGG A-3'), 0.25 μ L Hotstart Taq (Qiagen) enzyme, 5 μ L DNA template and free-water nuclease. The total volume is 50 μ L. The amplification process started with one cycle of pre-denaturation at 95°C for 7 minutes. It was followed by 40 cycles of the following steps: denaturation at 95°C for 45 minutes, followed by annealing 65°C for 45 seconds and extension at 72°C for 1 min. The gene polymorphism is determined using PCR-RFLP in 112 subjects, and the APOE gene was successfully amplified at 244 bp band, three alleles and six genotypes could be generated after excavation with HhaI endonuclease restriction (BioLabs).

Data Analysis

The statistical analysis was done to examine the relationship between APOE genotypes and alleles with MoCA-Ina questionnaire and its items (visuospatial, naming, attention, language, abstraction, memory delay, orientation).

RESULTS

MoCA-Ina questionnaire were conducted to determine the number of research participants who had with impaired cognitive function. The effect of APOE gene E4 on the cognitive function is different according to age.⁽¹⁷⁾

Table 1. Relationship between MoCA-Ina score and age

MoCA-Ina	Age (years), Frequency in % (n)		
	60 – 70	71 – 80	81 – 90
≥ 26	38.24 (13)	38.24 (13)	23.52 (8)
< 26	21.57 (11)	45.1 (23)	33.33 (17)

Fisher's Exact Test Exact: Sig. (2-sided) p 0.239

As shown in table 1; 85 participants divided into three groups: age group of 60-70, 71-80 and 81-90 years. The age group of 60 - 70 years: Percentage of participants with MoCA-Ina ≥ 26: 38.24% (13) and < 26: 21.57% (11). The age group of 71 - 80 years: Percentage of participants with MoCA-Ina ≥ 26: 38.24% (13) and < 26: 45.1% (23). For the age group of 81 - 90 years:

Percentage of participants with MoCA-Ina ≥ 26: 23.52% (8) and < 26: 33.33% (17). These results show an increase in the percentage of impaired cognitive function in the age group of 71-80 years, indicating the highest number of participants with mild cognitive impairments. There was an insignificant relationship between MoCA-Ina and age groups with p = 0.239.

Table 2. Relationship between age groups and MoCA-Ina items

Age Group	MoCA-Ina Item	P-Value (Mann-Whitney)
60-70 vs 71-80	Naming	0.035
	Orientation	0.036
71-80 vs 81-90	Naming	0.018
	Orientation	0.036
60-70 vs 81-90	Naming	0.000

Table 2 shows a significant decrease in the MoCA-Ina item naming when comparing the groups 60-70 vs. 71-80 years with p = 0.035, and the groups 71-80 vs. 81-90 years with p = 0.018. A significant decrease in the MoCA-Ina item orientation is also observed when comparing the groups the 60-70 vs. 71-80 years with p = 0.036, 71-80 vs. 81-90 years with p = 0.036 as well as 60-70 vs. 81-90 with p = 0.00.

Table 3. PCR-RFLP Analysis of APOE gene polymorphism

	Genotype Frequency, % (n)				Allele Frequency, % (n)		
	E2/E2	E3/E3	E3/E4	E4/E4	E2	E3	E4
Total = 112	8.9 (10)	33.9 (38)	41.1 (46)	16.1 (18)	8.9 (20)	54.5 (122)	36.6 (82)

Table 3 shows the APOE gene polymorphism by PCR-RFLP of all 112 participants. Results of shows the distribution of genotypes: E2/E2 8.9% (10), E3/E3 33.9% (38), E3/E4 41.1% (46), E4/E4 16.1% (18). The allele distribution were: E2 8.9% (20), E3 54.5% (122), E4 36.6% (82).

Table 4. Relationship MoCA-Ina score and APOE gene polymorphism

		Genotype frequency, % (n)				Allele frequency, % (n)		
		E2/E2	E3/E3	E3/E4	E4/E4	E2	E3	E4
MoCA-Ina Total= 85	≥26 (34)	8.8 (3)	20.6 (7)	47.1 (16)	23.5 (8)	8.8 (6)	44.1 (30)	47.1 (32)
	<26 (51)	13.7 (7)	27.5 (14)	43.1 (22)	15.7 (8)	13.7 (14)	49.0 (50)	37.3 (38)

Fisher's Exact Test Exact: Sig. (2-sided) p = 0.704

Table 4 shows the APOE gene polymorphism amongst 34 participants who obtained MoCA-Ina ≥ 26 : E2/E2 8.8%(3), E3/E3 20.6%(7), E3/E4 47.1%(16), E4/E4 23.5%(8). The 51 participants who obtained MoCA-Ina < 26 consist of: E2/E2 13.7%(7), E3/E3 27.5%(14), E3/E4 43.1%(22), E4/E4 15.7%(8).

Tabel 5 : Relationship APOE gene polymorphism and MoCA-Ina items

Genotype Comparison	MoCA-Ina Item	P-Value (Mann-Whitney)
E3/E4 vs E3/E3	Abstraction	0.019
E3/E4 vs E2/E2	Language	0.03
	Naming	0.048
E4/E4 vs E3/E3	Abstraction	0.028
E4/E4 vs E2/E2	Language	0.029

Table 5 shows a comparison between the APOE genotypes regarding items from MoCA-Ina in 85 participants. Analysis shows a significant relationship between APOE gene polymorphism and Abstraction: E3/E4 vs E3/E3 with $p = 0.019$, E4/E4 vs E3/E3 with $p = 0.028$, language: E3/E4 vs E2/E2 with $p = 0.03$, E4/E4 vs E2/E2 with $p = 0.029$, as well as the item naming: E3/E4 vs E2/E2 with $p = 0.048$. The allele E4 may be a contributing factor to the impairment of abstraction, language and naming abilities.

DISCUSSION

Memory seems to be of paramount importance as far as the cognitive decline is concerned, and other symptoms tend to receive less attention. This study shows that the age group 71-80 years is the most susceptible towards dementia with the highest percentage of individuals with impaired cognitive functions and the percentage decreases in the 81-90 years age group. This situation is not common as the risk of dementia increases with age. A significant drop in the item naming when comparing the age groups 60-70 vs. 71-80 with $p = 0.035$, and 71-80 vs. 81-90 with $p = 0.018$. A significant decrease is also observed for the item orientation when comparing the age groups 60-70 vs 71-80 with $p = 0.036$, 71-80 vs. 81-90 with $p = 0.036$ as well as 60-70 vs. 81-90 with $p = 0.00$.

The DNA analyzed using PCR RFLP of the 112 study participants found distribution of APOE gene

polymorphism: E2/E2 8.9%(10), E3/E3 33.9%(38), E3/E4 41.1%(46) and E4/E4 16.1%(18). The alleles distribution were: E2 8.9% (20), E3 54.5%(122), E4 36.6%(82). A study in Dong-gu Korea, found distribution of APOE gene polymorphism in the population: E2/E2 0.4%, E2/E3 10.7%, E2/E4 1.3%, E3/E3 71%, E3/E4 15.6% and E4/E4 1%. The alleles distribution were: E2 11.2%, E3 71%, E4 16.6%. Another Korean study in Namwon demonstrated the following genotype distribution: E2/E2 0.4%, E2/E3 10.2%, E2/E4 1.1%, E3/E3 72.6%, E3/E4 14.9% and E4/E4 0.9%. The alleles distribution as follows: E2 10.6%, E3 72.6%, E4 15.7%. The occurrence of the allele E4 in this study (Jakarta) is thus more-than-twice higher than that of the two above mentioned Korean studies. ⁽¹³⁾

The relationship between APOE gene polymorphism and the MoCA-Ina- item abstraction is as follows: E3/E4 vs. E3/E3 $p = 0.019$, E4/E4 vs. E3/E3 $p = 0.028$. Between APOE gene polymorphism and the item language: E3/E4 vs. E2/E2 $p = 0.03$, E4/E4 vs. E2/E2 $p = 0.029$. Between APOE gene polymorphism and the item naming: E3/E4 vs. E2/E2 $p = 0.048$. The E4 may be a potential factor contributing to the impairment of abstraction, language and naming abilities. Naming should receive attention as the dominant symptom of cognitive function decline. In a study by Coleman et al in Frontotemporal Dementia and Related Disorders patients with MoCA-Ina questionnaires, the sub-item naming and delay memory were most frequently detected in MCI / mild cognitive impairment patients. ⁽²¹⁾ This corresponds to the conclusion drawn by David A. Wolk et al. that APOE gene polymorphism has a significant effect on cognitive functions and brain anatomical abnormalities. ⁽¹⁴⁾

CONCLUSION

This study found that Decade seventh is the vulnerable group with the highest percentage with impaired cognitive functions. A high percentage of APOE allele E4 may be one of the causes of steep cognitive function decline and a higher risk for dementia. In determining the symptoms of dementia/ decline in cognitive function should not be fixated on memory loss but may be preceded by other symptoms, e.g., abstraction, language, orientation and especially naming.

Ethical Clearance - Taken from Hasanuddin University Ethics Committee, approval number: 130 /

H4.8.4.5.31 / PP36-KOMETIK / 2017.

Source of Funding- Scholarship from Medical Faculty, Tarumanagara University

Conflict of Interest- The author declares no conflict interest regard this research

REFERENCES

1. Wang X, Lopez OL, Sweet RA, Becker JT, DeKosky ST, Barnada MM, et al. Genetic determinants of disease progression in Alzheimer's disease. *Journal of Alzheimer's disease: JAD*. 2015;43(2):649.
2. Evans DA, Funkenstein HH, Albert MS, Scherr PA, Cook NR, Chown MJ, et al. Prevalence of Alzheimer's disease in a community population of older persons: higher than previously reported. *Jama*. 1989;262(18):2551-6.
3. Thalhauser CJ, Komarova NL. Alzheimer's disease: rapid and slow progression. *Journal of the Royal Society Interface*. 2011;rsif20110134.
4. Wang X, Wang H, Li H, Li T, Yu X. Frequency of the apolipoprotein E ϵ 4 allele in a memory clinic cohort in Beijing: a naturalistic descriptive study. 2014.
5. Chaudhry M, Hasnain S, Snitz B, Wang X, Rosenthal S, Demirci F, et al. Association of APOE polymorphisms and stressful life events with dementia in a Pakistani population. *Neuroscience letters*. 2014;570:42-6.
6. Jack C, Shiung M, Gunter J, O'Brien P, Weigand S, Knopman DS, et al. Comparison of different MRI brain atrophy rate measures with clinical disease progression in AD. *Neurology*. 2004;62(4):591-600.
7. Ridha BH, Barnes J, Bartlett JW, Godbolt A, Pepple T, Rossor MN, et al. Tracking atrophy progression in familial Alzheimer's disease: a serial MRI study. *The Lancet Neurology*. 2006;5(10):828-34.
8. Sluimer J, Vrenken H, Blankenstein M, Fox N, Scheltens P, Barkhof F, et al. Whole-brain atrophy rate in Alzheimer disease Identifying fast progressors. *Neurology*. 2008;70(19 Part 2):1836-41.
9. Mielke MM, Rosenberg P, Tschanz J, Cook L, Corcoran C, Hayden K, et al. Vascular factors predict rate of progression in Alzheimer disease. *Neurology*. 2007;69(19):1850-8.
10. Prolo P, Chiappelli F, Angeli A, Dovio A, Perotti P, Pautasso M, et al. Physiologic modulation of natural killer cell activity as an index of Alzheimer's disease progression. *Bioinformation*. 2007;1(9):363.
11. Rosyada A, Trihandini I. Determinan Komplikasi Kronik Diabetes Melitus pada Lanjut Usia. *Kesmas: National Public Health Journal*. 2013;7(9):395-402.
12. Yu B, Chen W, Wang R, Qi Q, Li K, Zhang W, et al. Association of apolipoprotein E polymorphism with maximal oxygen uptake after exercise training: a study of Chinese young adult. *Lipids Health Dis*. 2014;13:40.
13. Yun Y-W, Kweon S-S, Choi J-S, Rhee J, Lee Y-H, Nam H-S, et al. The APOE Polymorphism is Associated with C-reactive Protein Levels but Not with White Blood Cell Count: The Dong-gu Study and the Namwon Study. *Journal of Korean Medical Science*. 2014;30.
14. Wolk DA, Dickerson BC, Weiner M, Aiello M, Aisen P, Albert MS, et al. Apolipoprotein E (APOE) genotype has dissociable effects on memory and attentional-executive network function in Alzheimer's disease. *Proceedings of the National Academy of Sciences*. 2010;107(22):10256-61.
15. Poirier J, Bertrand P, Kogan S, Gauthier S, Davignon J, Bouthillier D. Apolipoprotein E polymorphism and Alzheimer's disease. *The Lancet*. 1993;342(8873):697-9.
16. Strittmatter WJ, Saunders AM, Schmechel D, Pericak-Vance M, Enghild J, Salvesen GS, et al. Apolipoprotein E: high-avidity binding to beta-amyloid and increased frequency of type 4 allele in late-onset familial Alzheimer disease. *Proceedings of the National Academy of Sciences*. 1993;90(5):1977-81.
17. Alfred T, Ben-Shlomo Y, Cooper R, Hardy R, Cooper C, Deary IJ, et al. Associations between APOE and low-density lipoprotein cholesterol genotypes and cognitive and physical capability: the HALCYon programme. *Age*. 2014;36(4):1-12.
18. Laws SM, Hone E, Gandy S, Martins RN. Expanding the association between the APOE gene and the risk of Alzheimer's disease: possible roles for APOE promoter polymorphisms and alterations

- in APOE transcription. *Journal of neurochemistry*. 2003;84(6):1215-36.
19. Reitz C, Jun G, Naj A, Rajbhandary R, Vardarajan BN, Wang L-S, et al. Variants in the ATP-binding cassette transporter (ABCA7), apolipoprotein E ϵ 4, and the risk of late-onset Alzheimer disease in African Americans. *Jama*. 2013;309(14):1483-92.
20. Shin M-H, Kweon S-S, Choi J-S, Lee Y-H, Nam H-S, Park K-S, et al. The effect of an APOE polymorphism on cognitive function depends on age. *Journal of neurology*. 2014;261(1):66-72.
21. Coleman KK, Coleman BL, MacKinley JD, Pasternak SH, Finger EC. Association between Montreal Cognitive Assessment sub-item scores and corresponding cognitive test performance in patients with frontotemporal dementia and related disorders. *Dementia and geriatric cognitive disorders*. 2017;43(3-4):170-9.