


# The Open Dermatology Journal





SJR

Scimago Journal &amp; Country Rank


Enter Journal Title, ISSN or Publisher Name

[Home](#)[Journal Rankings](#)[Country Rankings](#)[Viz Tools](#)[Help](#)[About Us](#)

# Open Dermatology Journal








**COUNTRY**[Netherlands](#)Universities and  
research institutions in  
Netherlands**SUBJECT AREA AND  
CATEGORY**[Medicine](#)  
[Dermatology](#)**PUBLISHER**[Bentham Science  
Publishers B.V.](#)**H-INDEX****9****PUBLICATION TYPE**[Journals](#)**ISSN**[18743722](#)**COVERAGE**[2010-2021](#)**INFORMATION**[Homepage](#)[How to publish in this  
journal](#)**SCOPE**

The Open Dermatology Journal is an open access online journal which publishes original research articles, reviews/mini-reviews, letters and guest edited thematic issues in all areas of experimental and clinical research in dermatology, skin disorders, cosmetic surgery, dermatitis, dermatopathology, dermatological surgery, immunodermatology, paediatric dermatology, teledermatology and reconstructive and cosmetic surgery. The Open Dermatology Journal, a peer-reviewed journal, is an important and reliable source of current information on developments in the field. The emphasis will be on publishing quality papers rapidly and freely available to researchers worldwide.

 [Join the conversation about this journal](#)[Quartiles](#)

FIND SIMILAR JOURNALS ?


<p>1 <b>Clinics in Dermatology</b> USA <b>68%</b> similarity</p>	<p>2 <b>International Journal of Women's Dermatology</b> NLD <b>65%</b> similarity</p>	<p>3 <b>American Journal of Clinical Dermatology</b> GBR <b>63%</b> similarity</p>	<p>4 <b>International Dermatology</b> GBR <b>6</b> s</p>
--	--	--	--



Show this widget in your own website

Just copy the code below and paste within your html code:

```
<a href="https://www.scimaç
```



**Open Dermatology Journal** ←


Q4 Dermatology  
best quartile

SJR 2021  
0.17

powered by scimagojr.com

**SCImago Graphica**

Explore, visually communicate and make sense of data with our [new data visualization tool](#).



Metrics based on Scopus® data as of April 2022



Loading comments...

Developed by:



Powered by:



Follow us on @ScimagoJR

Scimago Lab, Copyright 2007-2022. Data Source: Scopus®


**EST MODUS IN REBUS**  
Horatio (Satire 1,1,106)

[Edit Cookie Consent](#)

SPECIAL FEE WAIVER  
AND DISCOUNTS

GET BENTHAM OPEN  
MEMBERSHIP NOW!!

[Home](#) [About Journals](#) [Information for Authors/Reviewers](#) [Conferences](#) [Contact](#) [Help](#) [About Us](#) [Search](#)



## The Open Dermatology Journal

ISSN: 1874-3722 — Volume 14, 2020

- [Online Submission](#)
- [Submit Abstract Online](#)
- [Submit Issue Proposal](#)
- [Aims and Scope](#)
- [Abstracted / Indexed in](#)
- [View Published Contents](#)

Download Flyer



Announcing New  
Journal Website

- [New Journal Website](#)
- [Journal Home](#)
- [Editorial Board](#)
- [Meet the Executive Guest Editor\(s\)](#)
- [Board Recruitment Workflow](#)
- [Instructions for Authors](#)
- [Plagiarism Prevention](#)
- [Editorial Policies](#)
- [Publishing Ethics and Rectitude](#)
- [Quick Track Option](#)
- [Reviewer Guidelines](#)
- [Peer Review Workflow](#)
- [Guidelines for Guest Editors](#)
- [Publication Fee](#)
- [Publication Cycle - Process Flowchart](#)
- [Archiving Policies](#)

**Track Your Manuscript:**  
Enter Correct Manuscript  
Reference Number:

Submit Reference Number

### Endorsements

"Open access will revolutionize

### Browse Contents

- [Volume 14 - 2020](#)
- [Volume 13 - 2019](#)
- [Volume 12 - 2018](#)
- [Volume 11 - 2017](#)
- [Volume 10 - 2016](#)
- [Volume 9 - 2015](#)
- [Volume 8 - 2014](#)
- [Volume 7 - 2013](#)
- [Volume 6 - 2012](#)
- [Volume 5 - 2011](#)
- [Volume 4 - 2010](#)
- [Volume 3 - 2009](#)
- [Volume 2 - 2008](#)
- [Volume 1 - 2007](#)

Become an Editorial Board Member

[Register Here](#)

Become a Reviewer

[Register Here](#)

Become an Executive Guest Editor

[Register Here](#)

#### Editor-in-Chief:



**Craig G. Burkhart** (*Biography*)  
Department of Medicine  
University of Toledo  
Toledo, OH  
USA

#### Co-Editor:

**Cassian Sitaru** (*Biography*)  
Department of Dermatology  
Universitäts Klinikum Freiburg and Medical Faculty



Freiburg in Breisgau  
Germany

### Section Editor:



**(Current Frontier)**  
**Fabio Grizzi** (*Biography*)  
Department of Immunology and Inflammation  
Humanitas Research Hospital  
Milan  
Italy

### Editorial Board Members:



**B.J. Aneskievich** (*Biography*)  
Department of Pharmaceutical Sciences  
University of Connecticut  
Storrs, CT  
USA



**D. Bonamonte** (*Biography*)  
Department of Biomedical Science and Human  
Oncology  
University of Bari  
Bari  
Italy



**E. Berardesca** (*Biography*)  
Department of Clinical Dermatology  
San Gallicano Dermatological Institute  
Rome  
Italy



**R. Buquicchio** (*Biography*)  
Department of Dermatology  
University of Bari  
Bari  
Italy



**N. Chauhan** (*Biography*)  
Department of Otolaryngology - Head & Neck  
Surgery  
University of Toronto  
Toronto, ON  
Canada



**C.S. Chen** (*Biography*)  
Dermatology Service  
Memorial Sloan-Kettering Skin Cancer Center  
New York, NY  
USA



**D. Cho** (*Biography*)  
Department of Life Sciences  
Sookmyung Women's University  
Seoul  
S. Korea



**F. Cimino** (*Biography*)  
Department Farmaco-Biologico  
University of Messina  
Messina  
Italy



**A. Cowin** (*Biography*)  
Future Industries Institutes  
University of South Australia  
Adelaide  
Australia



**E. Errichetti** (*Biography*)  
Department of Experimental and Clinical  
Medicine  
University of Udine  
Udine  
Italy



**S. Eichmüller** (*Biography*)  
German Cancer Research Center  
GMP&T Cell Therapy Unit  
Heidelberg  
Germany



**J.Y. Fang** (*Biography*)  
Graduate Institute of Natural Products  
Chang Gung University  
Taoyuan  
Taiwan



**F. Furukawa** (*Biography*)  
Department of Dermatology  
Wakayama Medical University  
Wakayama  
Japan



**S. Garcovich** (*Biography*)  
Institute of Dermatology  
Catholic University of Sacred Heart  
Rome  
Italy



**C. Goh** (*Biography*)  
Department of Surgery  
National University of Singapore  
Singapore  
Singapore



**S. Gonzalez** (*Biography*)  
Department of Pathology  
Pontifical Catholic University of Chile  
Santiago  
Chile



**T. Hasegawa** (*Biography*)  
Department of Dermatology  
Juntendo University Graduate School of Medicine  
Tokyo  
Japan



**S.T. Hsieh** (*Biography*)  
Department of Neurology  
National Taiwan University Hospital  
Taipei  
Taiwan



**H. Ihn** (*Biography*)  
Department of Dermatology and Plastic and  
Reconstructive Surgery  
Kumamoto University  
Kumamoto  
Japan



**T. Kanekura** (*Biography*)  
Department of Dermatology  
Kagoshima University  
Kagoshima  
Japan



**G. Kroumpouzos** (*Biography*)  
Department of Dermatology  
Alpert Medical School of Brown University  
New York  
USA



**I. Kurokawa** (*Biography*)  
Department of Dermatology  
Meiwa Hospital  
Hyogo  
Japan



**A. Morita** (*Biography*)  
Department of Geriatric and Environmental  
Dermatology  
Nagoya City University Graduate School of  
Medical Sciences  
Nagoya  
Japan




**A.F. Nikkels** (*Biography*)  
Department of Dermatopathology  
University Hospital of Liège  
Liège  
Belgium



**T. Oberyszyn** (*Biography*)  
Department of Pathology  
The Ohio State University  
Columbus, OH  
USA



**N. Oyama** (*Biography*)  
Dermatology & Dermato-Allergology  
Matsuda General Hospital  
Ohno, Fukui  
Japan



**C.L. Tu** (*Biography*)  
VA Medical Center  
University of California  
San Francisco, CA  
USA



**M.T. Ventura** (*Biography*)  
Department of Internal Medicine, Immunology  
and Infectious Diseases  
University of Bari  
Bari  
Italy



**M. Vestita** (*Biography*)  
Department of Biomedical Sciences and Human  
Oncology  
University of Bari  
Bari  
Italy



**J.M. Weinberg** (*Biography*)  
Department of Dermatology  
St. Luke's-Roosevelt Hospital Center  
New York, NY  
USA



**C. Cerdeira** (*Biography*)  
Department of Dermatology  
University of Vigo  
Vigo  
Spain

**Associate Editorial Board Member:**



**J.A.M. Montanari** (*Biography*)  
Laboratory of Biomembranas  
National University of Quilmes  
Bernal  
Argentina



SPECIAL FEE WAIVER  
AND DISCOUNTS

GET BENTHAM OPEN  
MEMBERSHIP NOW!!

- [Home](#)
- [About Journals](#)
- [Information for Authors/Reviewers](#)
- [Conferences](#)
- [Contact](#)
- [Help](#)
- [About Us](#)
- [Search](#)



## The Open Dermatology Journal

ISSN: 1874-3722 — Volume 14, 2020

- [Online Submission](#)
- [Submit Abstract Online](#)
- [Submit Issue Proposal](#)
- [Aims and Scope](#)
- [Abstracted / Indexed in](#)
- [View Published Contents](#)



Announcing New  
Journal Website

- [New Journal Website](#)
- [Journal Home](#)
- [Editorial Board](#)
- [Meet the Executive Guest Editor\(s\)](#)
- [Board Recruitment Workflow](#)
- [Instructions for Authors](#)
- [Plagiarism Prevention](#)
- [Editorial Policies](#)
- [Publishing Ethics and Rectitude](#)
- [Quick Track Option](#)
- [Reviewer Guidelines](#)
- [Peer Review Workflow](#)
- [Guidelines for Guest Editors](#)
- [Publication Fee](#)
- [Publication Cycle - Process Flowchart](#)
- [Archiving Policies](#)

**Track Your Manuscript:**  
Enter Correct Manuscript  
Reference Number:

Submit Reference Number

### Endorsements

*"Open access will revolutionize 21<sup>st</sup> century knowledge work and accelerate the diffusion of ideas and evidence that support just in time learning and the evolution of thinking in a number of*

### Browse Contents

- [Volume 14 - 2020](#)
- [Volume 13 - 2019](#)
- [Volume 12 - 2018](#)
- [Volume 11 - 2017](#)
- [Volume 10 - 2016](#)
- [Volume 9 - 2015](#)
- [Volume 8 - 2014](#)
- [Volume 7 - 2013](#)
- [Volume 6 - 2012](#)
- [Volume 5 - 2011](#)
- [Volume 4 - 2010](#)
- [Volume 3 - 2009](#)
- [Volume 2 - 2008](#)
- [Volume 1 - 2007](#)

Become an Editorial Board Member <a href="#" style="background-color: white; color: #004a99; border-radius: 10px; padding: 2px 10px; display: inline-block;">Register Here</a>	Become a Reviewer <a href="#" style="background-color: white; color: #ff8c00; border-radius: 10px; padding: 2px 10px; display: inline-block;">Register Here</a>	Become an Executive Guest Editor <a href="#" style="background-color: white; color: #004a99; border-radius: 10px; padding: 2px 10px; display: inline-block;">Register Here</a>
---	--	---

## Volume 13 - 2019

### General Articles

CASE REPORT

**Bimodal Temporal Distribution of Herpes Explains Resistant Cases to Oral Antiviral Agents**

*The Open Dermatology Journal*, 2019, 13: 1-2

K.L. Gaishauer C.G. Burkhart

Electronic publication date: 31/1/2019


Collection year: 2019

Publisher Id: TODJ-13-1

DOI: 10.2174/1874372201913010001

[View Abstract](#)
[View Fulltext](#)
[Download PDF](#)
[Download ePub](#)





### Intrafamilial Diversity of Clinical Severity in Dominant Dystrophic Epidermolysis Bullosa: Case Series of Three Generations

*The Open Dermatology Journal*, 2019, 13: 3-7

Inne Arline Diana, Srie Prihianti Gondokaryono, Reiva Farah Dwiwana, July Iriani Rahardja, Yuri Yogya, Hendra Gunawan

**Electronic publication date:** 28/02/2019

**Collection year:** 2019

**Publisher Id:** TODJ-13-3

**DOI:** 10.2174/1874372201913010003

[View Abstract](#) [View Fulltext](#) [Download PDF](#) [Download ePub](#)



### REVIEW ARTICLE

#### Recent Insights On Diabetic Dermopathy

*The Open Dermatology Journal*, 2019, 13: 8-12

Rashmi Saxena Pal, Pranay Wal, Yogendra Pal, Ankita Wal

**Electronic publication date:** 28/02/2019

**Collection year:** 2019

**Publisher Id:** TODJ-13-8

**DOI:** 10.2174/1874372201913010008

[View Abstract](#) [View Fulltext](#) [Download PDF](#) [Download ePub](#)



### RESEARCH ARTICLE

#### Family Medicine Trainees' Knowledge about Topical Corticosteroids in Jeddah, Saudi Arabia

*The Open Dermatology Journal*, 2019, 13: 13-18

Amal A Kokandi

**Electronic publication date:** 30/04/2019

**Collection year:** 2019

**Publisher Id:** TODJ-13-13

**DOI:** 10.2174/1874372201913010013

[View Abstract](#) [View Fulltext](#) [Download PDF](#) [Download ePub](#)



### RESEARCH ARTICLE

#### Assessment of Vitamin D Status Among Cases with Pityriasis Alba

*The Open Dermatology Journal*, 2019, 13: 19-22

Wisam Majeed Kattoof, Ahmed Abdulhussein Kawen

**Electronic publication date:** 30/04/2019

**Collection year:** 2019

**Publisher Id:** TODJ-13-19

**DOI:** 10.2174/1874372201913010019

[View Abstract](#) [View Fulltext](#) [Download PDF](#) [Download ePub](#)



### CASE REPORT

#### Nummular Contact Eczema: Presentation of a Pediatric Case

*The Open Dermatology Journal*, 2019, 13: 23-26

D Bonamonte, A Filoni, G Gullo, M Vestita

**Electronic publication date:** 31/05/2019

**Collection year:** 2019

[View Abstract](#) [View Fulltext](#) [Download PDF](#) [Download ePub](#)



REVIEW ARTICLE

**The Benefits Of Occlusive Dressings In Wound Healing**

*The Open Dermatology Journal*, 2019, 13: 27-33


Sukmawati Tansil Tan, Nathasia Winarto, Ricky Dosan, Putri Bennya Aisyah

Electronic publication date: 31/05/2019

Publisher Id: TODJ-13-27

DOI: 10.2174/1874372201913010027

[View Abstract](#) [View Fulltext](#) [Download PDF](#) [Download ePub](#)



REVIEW ARTICLE

**Lessons From Epithelialization: The Reason Behind Moist Wound Environment**

*The Open Dermatology Journal*, 2019, 13: 34-40

Sukmawati Tansil Tan, Ricky Dosan

Electronic publication date: 31/07/2019

Collection year: 2019

Publisher Id: TODJ-13-34

DOI: 10.2174/1874372201913010034

[View Abstract](#) [View Fulltext](#) [Download PDF](#) [Download ePub](#)



REVIEW ARTICLE

**Current Review on Herbs for Derma Care**

*The Open Dermatology Journal*, 2019, 13: 41-46

Rashmi Saxena Pal, Yogendra Pal, Nikita Saraswat, Pranay Wal, Ankita Wal


Electronic publication date: 31/08/2019

Collection year: 2019

Publisher Id: TODJ-13-41

DOI: 10.2174/1874372201913010041

[View Abstract](#) [View Fulltext](#) [Download PDF](#) [Download ePub](#)



REVIEW ARTICLE

**Psoriasis Vulgaris: Relationship between Oral and Periodontal Conditions and Disease Severity**

*The Open Dermatology Journal*, 2019, 13: 47-54

Méndez Gaviria Ligia, Solis Leira, Rodríguez Constanza, Chila-Moreno Lorena, Buenahora María Rosa, Delgadillo Nathaly, Prieto Andrés, Castro Luis A, Romero-Sánchez Consuelo

Electronic publication date: 30/08/2019

Collection year: 2019

Publisher Id: TODJ-13-47

DOI: 10.2174/1874372201913010047

[View Abstract](#) [View Fulltext](#) [Download PDF](#) [Download ePub](#)



Madison Anzelc, Craig G. Burkhart

Electronic publication date: 15/11/2019  
Collection year: 2019

Publisher Id: TODJ-13-55  
DOI: 10.2174/1874372201913010055

[View Abstract](#) [View Fulltext](#) [Download PDF](#) [Download ePub](#)



#### CASE REPORT

##### Isolated Linear Discoid Lupus Erythematosus of Scalp Following Lines of Blaschko

*The Open Dermatology Journal*, 2019, 13: 58-60

Abdurrahman Almurayshid, Sulaiman Al Obaid

Electronic publication date: 20/12/2019  
Collection year: 2019

Publisher Id: TODJ-13-58  
DOI: 10.2174/1874372201913010058

[View Abstract](#) [View Fulltext](#) [Download PDF](#) [Download ePub](#)



#### RESEARCH ARTICLE

##### Clinical Aspects of Atopic Dermatitis of Children in Brazzaville, Congo.

*The Open Dermatology Journal*, 2019, 13: 61-67

Edith Sophie Bayonne-Kombo, Hébert Loubove, Yannichka Gloria Voumbo Mavaungou, Alphonse Gathsé

Electronic publication date: 31/12/2019  
Collection year: 2019

Publisher Id: TODJ-13-61  
DOI: 10.2174/1874372201913010061

[View Abstract](#) [View Fulltext](#) [Download PDF](#) [Download ePub](#)



#### REVIEW ARTICLE

##### Hydrogen Peroxide Use for Chemical Destruction in Seborrheic Keratosis: A Review

*The Open Dermatology Journal*, 2019, 13: 68-70

Darshana Balasubramaniam, Craig G. Burkhart

Electronic publication date: 31/12/2019  
Collection year: 2019

Publisher Id: TODJ-13-68  
DOI: 10.2174/1874372201913010068


[View Abstract](#) [View Fulltext](#) [Download PDF](#) [Download ePub](#)





# The Open Dermatology Journal

Content list available at: <https://opendermatologyjournal.com>



## REVIEW ARTICLE

### The Benefits Of Occlusive Dressings In Wound Healing

Sukmawati Tansil Tan\*, Nathasia Winarto, Ricky Dosan and Putri Bennya Aisyah

Department of Dermatovenereology, Faculty of Medicine, Tarumanagara University, Jakarta, Indonesia

**Abstract:** There are several types of wounds with their own healing properties. The latest innovation in wound management by using occlusive dressings can prevent infections, improve healing time and patient's comfort. Occlusive dressings are often used as an immediate wound hygiene control and also prevent blood loss until debridement is performed. They are used to protect wounds and surrounding tissue from pathogens and other harmful materials. A good cover depends on the condition around the wound, the person's skills, and the injury's nature. In this article, we provide an insight into the types of polymer materials used clinically in wound dressing and underlying mechanisms between the biomaterial dressings and the body tissue.

**Keywords:** Occlusive dressings, Moist wound environment, Wound healing, Chronic wounds, Acute wounds, Epidermis.

#### Article History

Received: February 23, 2019

Revised: April 22, 2019

Accepted: May 07, 2019

## 1. INTRODUCTION

In general, wounds can be classified into acute and chronic wounds. Acute wounds mostly tend to heal completely within 8–12 weeks with minimal scarring. Chronic wounds tend to reoccur and have a prolonged healing time that extends beyond 12 weeks. Wounds can also be classified according to skin layers and affected areas. Superficial wounds involve only the epidermal skin surface. Partial thickness wounds involve the epidermis, deeper dermal layers, blood vessels, sweat glands, and hair follicles. Full thickness wounds are caused by injuries that extend into the subcutaneous fat or deeper tissue along with the epidermis and dermal layers [1, 2].

In 1958, it was the first time known that moist environment for the wound could improve wound healing, when it was found that blisters healed faster if left in their original form [3]. Previously, honey paste, plant fiber, and animal fat were commonly used. Gauzes, lint, and cotton wool began to be used as wound dressings a few years later [4]. It was found in several studies that a polyethylene film made partial-thickness wounds of porcine heal approximately two times faster than the wound left open to the air. After that, many studies agreed about the benefits of moist wound environment for acute wound healing [3]. This has been found in acute but not chronic wounds because no animal model exists for chronic wounds. Due to excessive wound drainage from the traditional dressings, it becomes moistened and tends to become adherent

to the wound making it painful when removing. These traditional dressings fail to provide moist environment to the wound so they have been replaced by modern dressings with more advanced formulations [5]. Moist condition makes wound healing faster, relieves pain, prevents infection and contamination than acute wound treated with other methods [3, 6]. To prevent infections, occlusive dressings need to be changed regularly. Occlusive dressings are mostly produced in the form of a flat shape sheet. Some of them can be cut according to the size of the wound, but some of them also need secondary dressings to completely seal the wound [7].

## 2. WOUND HEALING

Wound healing is a complex process which is influenced by various mechanisms involving the coordinated interaction of blood cells, proteins, proteases, growth factors, and extracellular matrix components. Wound healing process can be divided into four phases (1) the coagulation and haemostasis phase; (2) inflammatory phase; (3) proliferative phase; and (4) maturational phase [8].

Hemostasis is initiated during the exposure of collagen during wound formation that activates the intrinsic and extrinsic clotting cascade. Injury to tissue causes a release of thromboxane A<sub>2</sub> and prostaglandin 2- $\alpha$  which leads to vasoconstriction. Extravasation of blood constituents assists in the formation of the blood clot. All of this process helps to limit hemorrhage and provides an initial extracellular matrix for cell migration.

After hemostasis is achieved, inflammatory phase begins. Capillary vasodilatation and leakage result secondary to local

\* Address correspondence to this author at the Department of Dermatovenereology, Tarumanagara University, Jakarta, Indonesia; Tel: +6287871111318; E-mail: sukmaskintreatment01@gmail.com

histamine release by the activated complement cascade. The increased blood flow and altered vascular permeability allow for the migration of inflammatory cells to the wound bed. Complement, neutrophil and macrophage play an important role in stimulating inflammatory cells, bacterial lysis and debris scavenging. In addition to direct phagocytosis of bacteria and foreign materials, macrophages secrete various enzymes and growth factors.

The proliferative phase is marked by epithelialization, angiogenesis, granulation tissue formation, and collagen deposition. Epithelialization occurs within hours after injury in wound healing and the normal layers of epidermis are restored typically in 3 days. The proliferative phase ends with granulation tissue formation.

The maturational phase is characterized by transition from the granulation tissue to scar formation. Around two weeks after injury, the wound undergoes contraction which results in a smaller amount of apparent scar tissue. Collagen deposition by fibroblasts continues for a prolonged period with a net increase in collagen deposition reaching after three weeks from tissue injury. The entire process is a dynamic continuum dictated by numerous growth factors and cells with an overlap of each of the three phases of wound healing to provide continued remodeling. The human wound has its maximal strength at one year [4, 9].

Promotion of these phases largely depends on the wound type, its associated pathological conditions and also the type of dressing material to achieve faster healing. Several factors can impair the whole process of wound healing. In general, these factors are classified as local and systemic. Local factors that affect wound healing are desiccation, infection, maceration, necrosis, oxygenation, pressure, trauma, and edema. (Checklist wound healing). Systemic factors such as aging, hormones, stress, and systemic diseases are essential in determining individual wound healing process [8, 10 - 14].

### 3. BASICS OF OCCLUSIVE DRESSINGS

Based on the wound type, suitable dressing material must be used. Dressing selection should be based on its ability to a) provide or maintain moist environment, b) enhance epidermal migration, c) promote angiogenesis and connective tissue synthesis, d) allow gas exchange between wounded tissue and environment, e) maintain appropriate tissue temperature to improve the blood flow to the wound bed and enhance epidermal migration, f) provide protection against bacterial infection, g) should be non-adherent to the wound and easy to remove after healing, h) must provide debridement action to enhance leucocytes migration and support the accumulation of enzyme, and i) must be sterile, non-toxic and non-allergic [15].

Modern wound dressings are usually based on synthetic polymers and are classified as passive, interactive and bioactive products. Passive products are non-occlusive, such as gauze and tulle dressings, used to cover the wound to restore its function underneath. Interactive dressings are semi-occlusive or occlusive, available in the form of films, foam, hydrogel and hydrocolloids. These dressings act as a barrier against penetration of bacteria to the wound environment [16 - 19].

Occlusive dressings increase about 40% of the process of

re-epithelialization of partial thickness wound [4, 20]. Studies where occlusive dressing leads to earlier epithelialization during wound healing have been proposed [20]. When the wound is closed with dressing, they are continuously exposed to proteinases, chemotactic, complement & growth factors, which is lost in the wound exposed. These dressings help in faster re-epithelialization, collagen synthesis, promote angiogenesis by creating hypoxia to the wound bed and decrease wound bed pH which leads to a decrease in the wound infection [21]. An increase in oxygen tension under occlusion has also been suggested as a beneficial effect of these dressings; however, hydrocolloids, which have healing rates as good as or better than those of polymer films, are actually oxygen impermeable, whereas the films are oxygen permeable.

Inflammatory phase and angiogenesis are enhanced in moist wound environment when compared to dry wound environment. Based on Winter *et al.* studies, dry wound tends to heal slowly with poor cosmesis compared with wound in a moist environment due to impaired reepithelialisation and scab formation [16, 19]. Fig. (1) describes the basic mechanism of occlusive dressings.

The number of endothelial and fibroblast cells increases in moist wound conditions as compared to wounds that are kept dry. Inflammatory phase cells such as neutrophils and macrophages increase in moist wound environment which can reduce scab or debris formation. By preventing wound desiccation, occlusion maintains an electrical gradient between wounded and non-wounded skin, which may stimulate epidermal cell migration.

Fluid in the wound improves wound healing by interacting with regenerating epithelium, dermal, and granulation tissue (ie, fibroblasts and endothelial cells). An increase in oxygen tension in combination with electrical stimulation has been shown to increase the expression of growth factor receptors on endothelial cells and fibroblast. Endothelial cells and fibroblast stimulated by platelet-derived growth factor increase collagen synthesis and promote angiogenesis. This mechanism plays an important role in preserving cosmesis. Inflammatory infiltrate develops from the dominant neutrophils to macrophages in mature wounds. Occlusion shortens the inflammatory healing phase and produces good cosmetic results. Fig. (2) describes the effects of occlusive dressings in wound healing process [4, 5].

It was thought that occlusion of wounds may lead to an increased risk of bacterial infection, but studies showed that wounds treated with dressings, which promote moist wound environment, are associated with a lower infection rate. Dermatitis has been reported after prolonged exposure to water, but such hydration induced changes to the epidermis can be reversed quickly if the underlying cause is removed. A wound dressing thus approaching ideal characteristics should conform to the site of the wound, offer alleviation of pain symptoms, promote faster wound-healing time and attempt to restore the patients' normal daily activities. Occlusive dressings protect the wound from both pathogenic invasion and further trauma. They act as barriers and prevent the entry of outside pathogens that might infect the wound and retard hea-




Fig. (1). Basic Mechanism of Occlusive Dressing.

ling. Since they conform to the body's contours, they are more likely to remain in place and offer protection from further trauma as well [4, 22 - 26].

**4. INDICATIONS AND CONTRAINDICATIONS FOR OCCLUSIVE DRESSINGS**

We summarized the indications and contraindications of occlusive dressings and also its beneficial effects.

Indications of occlusive dressings are [7]:

- Wound treatment. Occlusive dressings can be combined with antibiotics, gauze, sponge, hydrogels, and any other method to treat wounds
- Pressure and bleeding can often be immediately addressed
- Atopical ointments. It gives pressure to the ointment which improves absorption into the wound and prevents evaporation.
- Partial evisceration. Occlusive dressings are applied to protect the bowel until surgery.
- Sucking chest wounds. Wound and the puncture are covered and treated.

Contraindications of occlusive dressings are [7]:

- Maceration of the skin, where the skin feels tender, moist, and looks whiter than before, is an occlusive dressing's long-term risk.

- Pathogenic germs that exist in the infected area. Benefits of occlusive dressings [6]:

- Improve patient's comfort
- Lower the risk of infection
- Speed up wound healing

Several studies show the benefits of occlusive dressing for treating herpes virus skin infection. Based on histological analysis, it is known that skin injury caused by herpes virus is similar to partial-thickness skin wounds. Based on the study by Keegan *et al.*, occlusive dressing relieved pain and hypersensitivity caused by herpes zoster infection which may lead as an option to rule out the need for oral analgesics. A blinded control study by Lin *et al.* showed that the use of plain Tegaderm™ provided better pain relief. Lee *et al.* also showed the benefits of occlusive dressing for the management of herpes zoster by improving wound healing, relieving pain, improving patient's comfort and eschar debridement [27 - 29].

**5. TYPES OF OCCLUSIVE DRESSINGS**

Occlusive dressings work by maintaining moist wound environment which can shield the wound surface by preventing dry environment and trauma. Dry environment can prevent new epidermal cells migration to the wound surface. Variant types are: 1) semipermeable films, 2) hydrogels, 3) hydrocolloids and 4) alginates. Examples, functions, and features of each follow [6].




Fig. (2). Occlusive Dressing and Wound Healing Process.

### 5.1. Polymer Films

In 1960s, film dressings were introduced for wound management. These transparent semipermeable dressings are made of polyethylene or polyurethane with an adhesive coating on one side, are thin, highly elastomeric, elastic, permeable to moisture vapor and atmospheric gases but impermeable to liquid and bacteria [3]. Originally, these films are used for healing minor burns, superficial wounds, superficial decubitus ulcers, donor sites, blisters, abrasions, and cuts [6].

During the past years, wound dressing has been modernized which now mainly consists of synthetic polymers for wound management and can be classified as passive and interactive. Passive synthetic polymer dressings are non-occlusive, used for covering wound and help in restoring function under the polymer film. Examples of such passive synthetic polymers are gauze and tulle. Interactive synthetic

polymer dressings are occlusive or semi-occlusive which provide a barrier against bacterial penetration to the wound [30].

Polymer films trap exudates so the wound environment becomes moist [3, 4]. One main feature is that these films are impermeable to bacteria and liquid but are permeable to moisture vapor and air. The major drawback of polymer dressings is that exudates from wounds may accumulate underneath these dressings, which, due to increased pressure may cause a break in the environment maintained by the occlusive dressing [31, 32]. These products can be used for long-term bedridden patients acting as a shield to protect against skin friction in vulnerable areas, as a cover over topical products, and as intravenous dressings [3, 6].



**5.2. Polymer Foams**

Polyurethane foam film dressing is made of a thin microporous sheet of polyurethane foam that has been coated with a hydrophilic adhesive and bonded to a polyurethane film layer, which is good to avoid dehydration and bacterial infection. Foam film dressing has small pore size which increases the effectiveness of absorption or fluid retention. It also provides effectiveness in the prevention of bacterial infection and dehydration. Foam dressings attempt to rectify the lack of exudate absorbancy of occlusive dressing, without compromising on the moist environment needed for tissue repair [33 - 36]. Foams dressing does not cause maceration even if used for several days [4]

The major weakness of this type of dressing is that it needs secondary dressing like elastic bandage or a film to attach to the wound. Foam dressings are used for the wound that produced many exudates and deep wound or used under the compression bandage for chronic wound like venous ulcer [3, 4].

**5.3. Hydrogel Dressings**

Hydrogel contains about 96% water but does not dissolve in water and causes swelling if exposed to liquid solutions which can transmit moisture vapor. The high water content of hydrogels helps granulation tissues and epithelium in a moist environment. Hydrogel is made of polymers (polyethylene oxide, polyacrylamide, and polyvinylpyrrolidone) which are insoluble hydrophilic. Soft elastic property of hydrogels provides easy application and removal after the wound is healed without any damage. Their rate of absorbance is low, but high in capacity. For application, hydrogels need a secondary layer to attach to the wound, such as bandage or tape.<sup>3</sup> Hydrogel has soothing and cooling effect up to 5°C and maintains reduction in the cutaneous wounds temperature for up to 6 hours (may be augmented by refrigeration before use) which can alleviate pain and inflammation.

**Table 1. Wound Dressings Types and Purpose [1]**

Type	Purpose Use For
Polymeric hydrogels	Ulcers Chemotherapy peels Laser resurfacing
Polymeric foam	Burns Chronic wounds Laser resurfacing wounds Mohs surgery and wounds
Polymeric hydrocolloides	Average thickness wounds Burns Chronic Ulcer
Polymeric alginates	Surgical wounds Thickness burns Chronic ulcer High exudate wounds

The main disadvantage for using hydrogel is that it is easy to dry out which reduces their effectiveness and needs to be changed after 2 days [6]. Hydrogel is also a bad bacterial barrier.<sup>3</sup> Morgan has reported that except for infected and heavy drainage wounds, hydrogel dressings are suitable for all

four stages of wound healing. Difficulties in hydrogel dressings are that exudate accumulation leads to maceration and bacterial proliferation that produces foul smell in wounds. Besides, low mechanical strength of hydrogels makes it difficult to handle [36, 37]. Hydrogel is indicated for partial-thickness burns, dry chronic wounds, necrotic wounds, blisters and minor lacerations (Table 1) [3].

**5.4. Hydrocolloid Dressings**

Hydrocolloid is waterproof, semipermeable to vapor, opaque, occlusive, and absorbent. Hydrocolloid can be obtained synthetically or naturally [4]. It consists of two layers, inner colloidal layer and outer water-impermeable layer [3]. Hydrocolloids often use polysaccharides gel agents (sodium carboxymethyl cellulose), elastomeric compound, and are adhesive [4]. This type of dressing does not need secondary dressing and is impermeable to gases and moisture vapor [1]. Hydrocolloids are permeable to water vapor but impermeable to bacteria and also have the properties of debridement and absorb wound exudates [38]. They can melt so when they absorb exudate, but become viscous gel to stay in the wound when the dressings are removed. Hydrocolloids can be removed by normal saline [3].

The encapsulation properties of hydrocolloids create an opportunity to produce enhanced products with controlled drug (or any other appropriate substance) release due to the different type of hydrocolloid gels with different structural gelation. The amount of drug release can be controlled by optimization of the size of particles and the permeability of the gel membrane [39, 40].

Hydrocolloid dressings are designed on some wounds for prolonged periods (more than 1 week); this is useful in managing clean ulcers, but not when regular wound inspection is required. Thus, these dressings are probably more useful in preventing, rather than treating, infection within a wound [41]. The disadvantages of this dressing are the “gel and smell” phenomenon which may create an odorous residue when the exudate is mixed with adhesive when the dressing is removed.<sup>6</sup> In medical applications, the current market for hydrocolloid dressings offers their use in healing diabetic foot ulcers, chronic wound management, burn, and are also recommended for paediatric wound care management, as they do not cause pain on removal [3, 4].

**5.5. Alginates**

Alginate dressings are produced from the calcium and sodium salts of alginic acid which can be obtained from seaweed. Alginate dressings can be in the form of nonwoven mats, porous (foam) sheets or twisted staple fibers.<sup>3</sup> Alginate works by ion exchange reaction. The dressings would swell when wound exudates or sodium ions in tissue fluid are exchanged with alginates dressings of calcium ions. Alginate forms a gel in contact with the exudates. This property of alginate gels makes them very desirable as dressing for wounds that contain low to high exudates [42]. It would stay as gel for around one month if compared to hydrocolloids which degrade faster [3, 4].

**Table 2. Occlusive Dressings Benefits<sup>1</sup>**

	Polymer Foams	Hydrogels	Polymer Films	Alginates	Hydrocolloids
Absorbent	+	+	-	+	±
Transparent	-	±	+	-	-
Suited for exudative wounds	+	±	-	+	-
Permeable to water vapor	+	+	+	+	-
Permeable to oxygen	+	+	+	+	-
Prevents bacterial entry	-	-	-	-	+
Adherent to nonwounded skin	-	-	+	-	+

One of the main reasons for using alginates in wound dressings is their haemostatic ability; therefore, alginate dressings can be used for wounds that are bleeding. The coagulation effects of zinc and calcium alginate dressings have been compared with non-alginate dressings. Zinc containing alginates have the best haemostatic ability. Alginate dressings have an important property of gel formation which helps in reducing the pain during removal and changing of the bandage from the wound site. Alginate gels can also be used for the delivery of drugs having low molecular weight. The partially oxidized alginate gels are known to be used for controlled and localized delivery of antineoplastic agents [43, 44].

Alginate dressings have applications in tissue regeneration and bio-engineering fields. Depending on composition, sodium alginate has been identified as a substrate for cell proliferation. This opens up new possibilities for tissue regeneration in skin scaffolds as well [45]. Even though some studies have reported that alginate inhibits keratinocytes migration, Thomas *et al.*, have reported that alginates accelerate healing process by activating macrophages to produce TNF- $\alpha$  which initiates inflammatory signals [46]. In dry condition, alginate dressing removal may damage the newly formed epithelium around the wound. Alginate dressings are suitable for moderate to heavy drainage wounds and not suggested for dry wound, third degree burn wound and severe wounds with exposed bone. Like polymer foam and hydrogel, alginate also needs a secondary dressing to be attached (Table 2) [3].

## CONCLUSION

Wound dressings have experienced development to more active dressings which create moist wound environment. The ideal characteristics of an occlusive dressings are to maintain moist environment, shorten and accelerate wound healing process such as; enhance epidermal migration, promote angiogenesis and connective tissue synthesis, allow gas exchange, improve blood flow to the wound bed and enhance epidermal migration. Moreover, they provide protection against bacterial infection, are non-adherent to the wound providing debridement action, are sterile, non-toxic and non-allergic. Occlusive dressings are not suitable to treat wounds with impaired circulation such as burn ulcers, pressure ulcers, or other heavy trauma.

## CONSENT FOR PUBLICATION

Written consent was obtained from the patient for the surgery and the publication of her data.

## AVAILABILITY OF DATA AND MATERIALS

Not applicable.

## FUNDING

None.

## CONFLICT OF INTEREST

The authors declare no conflict of interest, financial or otherwise.

## ACKNOWLEDGMENTS

Declared none.

## REFERENCES

- [1] Bryant R, Nix D. Acute and chronic wounds. Amsterdam: Elsevier Health Sciences 2006.
- [2] Bolton L, van Rijswijk L. Wound dressings: Meeting clinical and biological needs. *Dermatol Nurs* 1991; 3(3): 146-61. [PMID: 1828677]
- [3] Helfman T, Ovington L, Falanga V. Occlusive dressings and wound healing. *Clin Dermatol* 1994; 12(1): 121-7. [http://dx.doi.org/10.1016/0738-081X(94)90262-3] [PMID: 8180934]
- [4] Mir M, Ali MN, Barakullah A, *et al.* Synthetic polymeric biomaterials for wound healing: a review. *Prog Biomater* 2018; 7(1): 1-21. [http://dx.doi.org/10.1007/s40204-018-0083-4] [PMID: 29446015]
- [5] Boateng JS, Matthews KH, Stevens HNE, Eccleston GM. Wound healing dressings and drug delivery systems: A review. *J Pharm Sci* 2008; 97(8): 2892-923. [http://dx.doi.org/10.1002/jps.21210] [PMID: 17963217]
- [6] Scot B. MS, ATC, PA-S. Wound Management: The Occlusive Dressing. *J Athl Train* 1995; 30(2): 143-46.
- [7] WoundSource. What is an Occlusive Dressing?. 2016.
- [8] Dhivya S, Padma VV, Santhini E. Wound dressings - A review. *Biomedicine (Taipei)* 2015; 5(4): 22. [http://dx.doi.org/10.7603/s40681-015-0022-9] [PMID: 26615539]
- [9] Sinno H, Prakash S. Complements and the wound healing cascade: an updated review. *Plast Surg Int* 2013; 2013(146764)146764 [http://dx.doi.org/10.1155/2013/146764] [PMID: 23984063]
- [10] Guo S, Dipietro LA. Factors affecting wound healing. *J Dent Res* 2010; 89(3): 219-29. [http://dx.doi.org/10.1177/0022034509359125] [PMID: 20139336]
- [11] Godbout JP, Glaser R. Stress-induced immune dysregulation: implications for wound healing, infectious disease and cancer. *J Neuroimmune Pharmacol* 2006; 1(4): 421-7. [http://dx.doi.org/10.1007/s11481-006-9036-0] [PMID: 18040814]
- [12] Brem H, Tomic-Canic M. Cellular and molecular basis of wound healing in diabetes. *J Clin Invest* 2007; 117(5): 1219-22. [http://dx.doi.org/10.1172/JCI32169] [PMID: 17476353]
- [13] Wilson JA, Clark JJ. Obesity: impediment to postsurgical wound healing. *Adv Skin Wound Care* 2004; 17(8): 426-35. [http://dx.doi.org/10.1097/00129334-200410000-00013] [PMID: 1549 2679]
- [14] Thomas Hess C. Checklist for factors affecting wound healing. *Adv Skin Wound Care* 2011; 24(4): 192. [http://dx.doi.org/10.1097/01.ASW.0000396300.04173.ec] [PMID: 2142 2844]

- [15] Degreef HJ. How to heal a wound fast. *Dermatol Clin* 1998; 16(2): 365-75. [http://dx.doi.org/10.1016/S0733-8635(05)70019-X] [PMID: 9589 210]
- [16] Hunt TK, Hopf H, Hussain Z. Physiology of wound healing. *Adv Skin Wound Care* 2000; 13(2)(Suppl.): 6-11. [PMID: 11074996]
- [17] Rivera AE, Spencer JM. Clinical aspects of full-thickness wound healing. *Clin Dermatol* 2007; 25(1): 39-48. [http://dx.doi.org/10.1016/j.clindermatol.2006.10.001] [PMID: 17276 200]
- [18] Strecker-McGraw MK, Jones TR, Baer DG. Soft tissue wounds and principles of healing. *Emerg Med Clin North Am* 2007; 25(1): 1-22. [http://dx.doi.org/10.1016/j.emc.2006.12.002] [PMID: 17400070]
- [19] Winter GD. Effect of air exposure and occlusion on experimental human skin wounds. *Nature* 1963; 200: 378-9. [http://dx.doi.org/10.1038/200378a0] [PMID: 14087905]
- [20] Nayeri F. Occlusive bandaging of wounds with decreased circulation promotes growth of anaerobic bacteria and necrosis: case report. *BMC Res Notes* 2016; 9: 394. [http://dx.doi.org/10.1186/s13104-016-2205-1] [PMID: 27502024]
- [21] Sarabahi S. Recent advances in topical wound care. *Indian J Plast Surg* 2012; 45(2): 379-87. [http://dx.doi.org/10.4103/0970-0358.101321] [PMID: 23162238]
- [22] Winter GD, Scales JT. Effect of air drying and dressings on the surface of a wound. *Nature* 1963; 197: 91-2. [http://dx.doi.org/10.1038/197091b0] [PMID: 14001241]
- [23] Ousey K, Cutting KF, Rogers AA, Rippon MG. The importance of hydration in wound healing: reinvigorating the clinical perspective. *J Wound Care* 2016; 25(3): 122-130, 124-130. [http://dx.doi.org/10.12968/jowc.2016.25.3.122] [PMID: 26947692]
- [24] Björklund S, Ruzgas T, Nowacka A, *et al.* Skin membrane electrical impedance properties under the influence of a varying water gradient. *Biophys J* 2013; 104(12): 2639-50. [http://dx.doi.org/10.1016/j.bpj.2013.05.008] [PMID: 23790372]
- [25] Thomas S. 2008 The role of dressings in the treatment of moisture-related skin damage. *World Wide Wounds* [February 2016]; http://bit.ly/1wRQThs
- [26] Rheinecker SB. Wound management: the occlusive dressing. *J Athl Train* 1995; 30(2): 143-6. [PMID: 16558325]
- [27] Keegan DA. Reducing pain in acute herpes zoster with plain occlusive dressings: A case report. *J Med Case Reports* 2015; 9: 89. [http://dx.doi.org/10.1186/s13256-015-0560-5] [PMID: 25907451]
- [28] Lin PL, Fan SZ, Huang CH, *et al.* Analgesic effect of lidocaine patch 5% in the treatment of acute herpes zoster: A double-blind and vehicle-controlled study. *Reg Anesth Pain Med* 2008; 33(4): 320-5. [http://dx.doi.org/10.1097/00115550-200807000-00006] [PMID: 1867 5742]
- [29] Lee SK. Notes on Practice: Healing Shingles with Moist Occlusive Dressings. *Wound Management & Prevention* 2002; 40(3)
- [30] Dhivya S, Padma VV, Santhini E. Wound dressings - A review. *Biomedicine (Taipei)* 2015; 5(4): 22. [http://dx.doi.org/10.7603/s40681-015-0022-9] [PMID: 26615539]
- [31] Jenks M, Craig J, Green W, Hewitt N, Arber M, Sims A. Tegaderm CHG IV securement dressing for central venous and arterial catheter insertion sites: A NICE medical technology guidance. *Appl Health Econ Health Policy* 2016; 14(2): 135-49. [http://dx.doi.org/10.1007/s40258-015-0202-5] [PMID: 26458938]
- [32] Khil MS, Cha DI, Kim HY, Kim IS, Bhattarai N. Electrospun nanofibrous polyurethane membrane as wound dressing. *J Biomed Mater Res B Appl Biomater* 2003; 67(2): 675-9. [http://dx.doi.org/10.1002/jbm.b.10058] [PMID: 14598393]
- [33] Hinrichs WL, Lommen EJ, Wildevuur CR, Feijen J. Fabrication and characterization of an asymmetric polyurethane membrane for use as a wound dressing. *J Appl Biomater* 1992; 3(4): 287-303. [http://dx.doi.org/10.1002/jab.770030408] [PMID: 10147998]
- [34] Helfman T, Ovington L, Falanga V. Occlusive dressings and wound healing. *Clin Dermatol* 1994; 12(1): 121-7. [http://dx.doi.org/10.1016/0738-081X(94)90262-3] [PMID: 8180934]
- [35] Zahedi P, Rezaeioam O, Ranaei-Siadat S, Jafari S, Supaphol P. A review on wound dressings with an emphasis on electrospun nanofibrous polymeric bandages. *Polym Adv Technol* 2010; 21: 77-95.
- [36] Thomson T. Foam Composite. US Patent 7048966, 2006.
- [37] Martin L, Wilson CG, Koosha F, *et al.* The release of model macromolecules may be controlled by the hydrophobicity of palmitoyl glycol chitosan hydrogels. *J Control Release* 2002; 80(1-3): 87-100. [http://dx.doi.org/10.1016/S0168-3659(02)00005-6] [PMID: 11943 390]
- [38] Thomas S, Loveless PA. A comparative study of twelve hydrocolloid dressings. *World Wide Wounds* 1997; 1: 1-12.
- [39] Bramhill J, Ross S, Ross G. Bioactive nanocomposites for tissue repair and regeneration: A review. *Int J Environ Res Public Health* 2017; 14(1): 66. [http://dx.doi.org/10.3390/ijerph14010066] [PMID: 28085054]
- [40] Das S, Baker AB. Biomaterials and nanotherapeutics for enhancing skin wound healing. *Front Bioeng Biotechnol* 2016; 4: 82. [http://dx.doi.org/10.3389/fbioe.2016.00082] [PMID: 27843895]
- [41] Frykberg RG, Banks J. Challenges in the treatment of chronic wounds. *Adv Wound Care (New Rochelle)* 2015; 4(9): 560-82. [http://dx.doi.org/10.1089/wound.2015.0635] [PMID: 26339534]
- [42] Dabiri G, Damstetter E, Phillips T. Choosing a wound dressing based on common wound characteristics. *Adv Wound Care (New Rochelle)* 2016; 5(1): 32-41. [http://dx.doi.org/10.1089/wound.2014.0586] [PMID: 26858913]
- [43] Segal HC, Hunt BJ, Gilding K. The effects of alginate and non-alginate wound dressings on blood coagulation and platelet activation. *J Biomater Appl* 1998; 12(3): 249-57. [http://dx.doi.org/10.1177/088532829801200305] [PMID: 9493071]
- [44] Paul W, Sharma CP. Chitosan and alginate wound dressings. *Trends Biomater Artif Organs* 2004; 18(1): 18-23.
- [45] Wang L, Shelton RM, Cooper PR, Lawson M, Triffitt JT, Barralet JE. Evaluation of sodium alginate for bone marrow cell tissue engineering. *Biomaterials* 2003; 24(20): 3475-81. [http://dx.doi.org/10.1016/S0142-9612(03)00167-4] [PMID: 12809 776]
- [46] Thomas A, Harding KG, Moore K. Alginates from wound dressings activate human macrophages to secrete tumour necrosis factor-alpha. *Biomaterials* 2000; 21(17): 1797-802. [http://dx.doi.org/10.1016/S0142-9612(00)00072-7] [PMID: 10905 462]